

INTRAUTERINE INSEMINATION (IUI)

INSIDE:

- What to expect from fertility medications
- IUI explained step by step
- Coping with stress



ABOUT THIS BOOKLET

This series of booklets has been developed and written with the support of leading fertility clinics across Australia, and Access Australia – a national patient advocacy organisation that provides numerous services for people having difficulty conceiving. We also acknowledge the many people who spoke openly about their own experiences with assisted conception in order to help others experiencing a similar journey. Merck thanks the many individuals, couples and Australian healthcare professionals, including fertility specialists, specialist nurses and psychologists who shared their knowledge and expertise during the production of these booklets. Merck acknowledges the contribution of Dr Gareth Weston in the preparation of this booklet.

Important notice: The information provided in this booklet does not replace any of the information or advice provided by a medical practitioner and other members of your healthcare team. Your doctor will determine the best medications and course of action for you based on your requirements and circumstances.

Prescription medicines have benefits and risks. Use all prescribed medicines strictly as directed by your doctor and raise any questions or concerns with them before, during or after using them. If you experience side effects consult your doctor.

Medication availability and funding criteria may differ between Australia and NZ.



CONTENTS

About this booklet	2
Introduction	4
What is intrauterine insemination (IUI)?	4
What are the benefits?	5
When is IUI not suitable?	5
Reasons for using IUI	6
How is IUI performed?	7
Ovulation.....	7
Monitoring ovulation	8
Semen collection	9
Sperm preparation	10
Insertion	10
Success rates	11
Coping with stress	12
Support organisations	13

INTRODUCTION

Having trouble falling pregnant comes as a surprise to most women and men. Many people assume that pregnancy will follow immediately after birth control is discontinued. In reality, up to one in six couples worldwide have difficulty conceiving during the first 12 months of trying.¹

There are many treatments available that may be able to help you to become pregnant.

Intrauterine insemination (IUI) is a common procedure that has been used for many years to help deliver male sperm closer to the female egg.

WHAT IS INTRAUTERINE INSEMINATION (IUI)?²

You may have heard the term **artificial insemination (AI)**. This is the name given to a procedure where sperm are placed into the female reproductive system by a means other than intercourse. **IUI** is the most common form of AI used and involves placing sperm into the female's uterus through an assisted medical process. As it is a relatively low-tech solution to infertility problems, IUI is usually one of the first techniques used to assist a couple who is having difficulty becoming pregnant.

In the procedure, warmed and 'washed' (treated) sperm are introduced into the woman's uterus through a flexible tube. Sperm can be provided by the woman's husband or partner (artificial insemination by husband – AIH) or sperm provided by a known or anonymous sperm donor (artificial insemination by donor – AID or DI). The procedure is done around the time of ovulation to give the best chance of conception. Hormonal (fertility) medications might be used in conjunction with the treatment to enhance conditions for a pregnancy, either by increasing the number of follicles and eggs produced with a cycle, or triggering release of the egg(s) at a precisely known time to coordinate better with insemination.

WHAT ARE THE BENEFITS OF IUI?³

During normal intercourse, only relatively small numbers of sperm make it to the woman's uterus and into the fallopian tubes where fertilisation takes place. IUI inserts large amounts of the best performing sperm directly into a woman's uterus thus increasing the chance of fertilisation.



WHEN IS IUI NOT SUITABLE?⁵

IUI is not effective when there is:

- bilateral/blocked or damaged fallopian tubes
- ovarian failure (menopause)
- severe male factor infertility, i.e. no sperm, very low sperm count, poor sperm motility (movement), sperm defects; in this case, donor sperm may be an option
- severe endometriosis.

REASONS FOR USING IUI³

IUI is mainly used when timed intercourse or hormonal medications alone have not worked, or if there are 'mild' sperm abnormalities, such as poor motility (i.e. the ability of the sperm to move). It is also used in conjunction with donor sperm by same sex female couples or by women who don't have a partner.



In addition, IUI can be used to overcome fertility due to the following conditions:

- **mild endometriosis:** occurs when the tissue that normally lines the inside of the uterus grows in other places where it doesn't belong, such as on the ovaries and fallopian tubes
- **mucus 'hostility':** may arise as a result of a vaginal infection or the presence of antisperm antibodies in the mucus
- **ovulation problems:** including irregular or absent ovulation often caused by a deficiency in one of the controlling hormones
- **ejaculation problems:** due to psychological problems such as impotence (inability to get and maintain an erection), or anatomic problems of the penis, such as paraplegia
- **retrograde ejaculation:** the semen goes backward into the bladder instead of coming out of the penis
- **unexplained infertility:** defined as not being able to conceive after one year of unprotected intercourse, even though the cycle is normal, semen is normal, internal examinations are normal and there is normal sperm-mucus penetration
- **physical problems with sexual intercourse:** vaginismus is an involuntary clamping of the vaginal muscles, which may prevent some women from having frequent intercourse.

HOW IS IUI PERFORMED?



Ovulation³

There are different ways your doctor may choose to conduct IUI depending on your individual situation.

IUI done without hormonal medication

If the woman has a regular menstrual cycle, IUI will be performed about Day 12 to Day 15 of the cycle when ovulation – release of the egg – is taking place. It is also suitable for those who cannot have natural intercourse (e.g. spinal cord injuries). You will be asked to work out when ovulation will occur by tracking basal body temperature (your temperature increases during ovulation) and changes in vaginal mucus, or by using ovulation kits. Alternatively, you may be monitored through regular ultrasounds or blood tests administered by your medical team or clinic.

IUI done in conjunction with hormonal medications

Depending on your particular situation, your doctor may recommend that you take hormonal medication (also known as ‘fertility drugs’) to help stimulate ovulation. Your doctor may refer to this part of treatment as ‘ovulation induction’, ‘ovarian stimulation’ or ‘stimulated cycle’.

THE RISK OF MULTIPLE PREGNANCIES³

Fertility medications can cause more than one follicle (egg) to develop and this is why your body will be monitored closely through blood tests and ultrasounds while you are taking them.

For those having trouble becoming pregnant, having twins may seem like a blessing, but complications, such as miscarriage and low birth weight, are much more common in twins than single pregnancies.

Monitoring ovulation

Throughout this first stage of fertility treatment, your response to follicle stimulating hormone (FSH) will be carefully monitored for ovarian hyperstimulation syndrome (OHSS) – see box below – and to gain a clearer picture of what is happening to the follicles so the right timing and dose of fertility medication can be determined. This monitoring will be done through regular ultrasounds, blood tests and urine tests.



OVARIAN HYPERSTIMULATION SYNDROME

Ovarian hyperstimulation syndrome (OHSS) is a potentially life-threatening medical condition which may occur when your ovaries have been overly stimulated by various fertility medications. The ovaries may increase in size and produce large amounts of fluid. It is characterised by pain and bloating in your abdomen and if severe, can cause problems with breathing or urination. Contact a member of your healthcare team immediately if you believe you have any of these symptoms.

Semen collection⁶

Artificial insemination by partner

On the day of the insemination, the male partner will be required to produce a sample of semen by ejaculating into a sterile container. Two to three days' abstinence from intercourse/masturbation is preferred prior to the sample collection day. Clinics often provide a room so that this sample may be produced in private, but some men prefer to collect the semen at home and deliver it to the clinic. As it is being used that day it must not be frozen or refrigerated, and it needs to arrive promptly at the clinic – within a couple of hours.

Donor insemination (DI)

IUI can also be done using donor sperm, either from an anonymous or a known sperm donor (known as DI or donor insemination). Insemination with donor sperm is used when there is no male partner or the male partner does not produce sperm, when the sperm are of very poor quality or if there is a high risk of passing on genetic diseases.

Sperm are frozen ahead of time, quarantined for 3–6 months, and screened for sexually transmitted diseases (e.g. HIV/AIDS/Hep B/Hep C) and common genetic disorders. The semen selected for a couple closely matches, as much as possible, the male partner's characteristics, e.g. eye and hair colour, height and build.

Many clinics will allow couples to reserve semen for a subsequent pregnancy, so that if they have a child from insemination, they can try for another child with the same genetic characteristics (i.e. from the same donor).

How might you feel?

Using donor sperm

Before you agree to use donor sperm, it is important for you to explore how you truly feel about it. For example, the child created will be genetically related to only one of you. For the male partner, this may result in feeling like your masculinity or even your relationship is threatened and that you are no longer part of the conception/pregnancy process. On the contrary, you will be there from the point of insemination right through to raising your child. Providing sperm does not automatically mean you will be a good father. Being a parent is about passing on your values, love, wisdom and experience.

It is a good idea to discuss your emotions and concerns with a counsellor available through your fertility clinic or as recommended by your doctor. **AccessAustralia** and the **Australian Donor Conception Network** have many resources on the issues relating to donor insemination (see contact details at the back of this booklet).


Sperm preparation³

The semen is prepared in a laboratory for the next stage. In the woman's body, the cervix acts as a filter for the sperm, so only the motile sperm pass through. In the same way, during the IUI procedure the semen is 'washed' and filtered, removing any mucus and non-motile sperm. In other words, a concentrated solution containing the most active sperm is inserted. Unwashed sperm must not be placed in the uterus as severe allergic reactions can occur.

The samples are thawed and the most active sperm are separated as required. Additional 'straws' or vials of sperm can be thawed if required to give a suitable insertion sample.

Insertion³

On the day of ovulation, sperm insertion will take place with fresh or thawed sperm. For those using fresh sperm, you will be asked to return to the clinic one to two hours after the delivery of the sperm sample for the insertion process. This simple and low-tech procedure is done without anaesthetic and is rather like having a pap smear, with a similar level of discomfort involved. An instrument called a speculum is inserted into the vagina so that the cervix (neck of the uterus) can be seen. A narrow tube attached to a syringe is gently pushed through the cervix into the uterus. Sometimes when the cervix is hard to reach a longer tube may be



Adapted from Gardner DK *et al*, 2004.⁴

attached to the end of the syringe. The sperm are then injected into the uterus. It is usually not painful but occasionally some mild cramping or discomfort may occur. You will be asked to remain lying down for 10 to 15 minutes and then you will be able to resume your normal routine.

If you are on a FSH/hCG/IUI cycle, you may need a booster injection of hCG – normally done at home – around seven days after insertion.

What can help?

Surviving the two-week wait

After your IUI, it takes around two weeks for pregnancy test results to be accurate. This 'two-week wait' – the time before your expected period – is understandably a time of high anxiety, worry and frustration for women trying to conceive. Here are some 'survival' tips to help you get through this time:

- try not to obsess about symptoms of pregnancy – feeling pregnant does not always mean that you are. Your doctor may prescribe medication after your treatment with side effects that resemble pregnancy symptoms
- keep busy – this may mean working more, or planning meaningful or fun distractions
- allow yourself a limited amount of time each day to think about pregnancy, write down your thoughts, search information online or discuss it with your partner or supportive friends/family members
- try some relaxation techniques such as breathing exercises or meditation

SUCCESS RATES

The success rates for IUI will vary depending on a number of factors including the number of cycle attempts and maternal age. You should discuss success rates with your doctor.

COPING WITH STRESS

Talk to your partner

Infertility is a couple's problem not an individual's. By asking for and relying on the support of your partner and by communicating openly with them throughout the evaluation, diagnosis and treatment phases, you may find that your relationship grows stronger.



The value of a wide support network

Discussing fertility issues may be uncomfortable, but expressing how you feel may help you release your stress. It's important that you reach out for support.

Infertility is a sensitive subject and many people may not know how to react. Guide the conversation and help them avoid topics that may be hurtful or make you feel uncomfortable. Let your friends know how they can support you.

If you feel in need of more emotional support than your partner can give, but don't want to share everything with a friend or family member, your fertility clinic usually offers the services of a counsellor.

SUPPORT ORGANISATIONS

AUSTRALIA

Access Australia

www.access.org.au

Ph: 1800 888 896

Email: info@access.org.au

Access Australia is a national patient advocacy organisation, which provides numerous services and resources for people having difficulty conceiving. Its services include:

- fact sheets, newsletters and personal stories
- putting you in contact by phone or email with others sharing a similar infertility experience
- a register of infertility self-help groups
- listing of infertility clinics accredited by the Reproductive Technology Accreditation Committee (RTAC)
- listing of professional infertility counsellors across Australia
- lobbying governments for equal access to affordable, quality assisted conception treatment.

Australian Donor Conception Network

www.australiandonorconceptionnetwork.org

Email: donorconceptionnetwork@gmail.com

The Australian Donor Conception Network is an organisation run by volunteers. Its members include those who are considering or using donor sperm, egg or embryo, those who already have children conceived on donor programs, adult donor offspring and donors. It offers a social events for members to share their experiences, Facebook groups to enable members to stay in touch, informative emails, a library of books and links to other helpful organisations.

Endometriosis Care Centre of Australia

www.ecca.com.au

Formed by a group of health specialists, this organisation provides patient information and a state by state 'find a specialist' search engine on its website.

Endometriosis Australia

www.endometriosisaustralia.org

Email: admin@endoaustralia.org

Provides information on state contacts.

Endometriosis Australia endeavours to increase recognition of endometriosis, provide endometriosis education programs, and help fund endometriosis research. They strive to build strong relationships with existing endometriosis support networks throughout the country.

SANDS (miscarriage, stillbirth & newborn death support)

www.sands.org.au

Ph: 1300 072 637

Email: support@sands.org.au

SANDS is a self-help support group comprised of parents who have experienced the death of a baby through miscarriage, stillbirth, or shortly after birth. It provides 24-hour telephone support, information resources, monthly support meetings, name-giving certificates and other support.

SUPPORT ORGANISATIONS

NEW ZEALAND

Fertility NZ

www.fertilitynz.org.nz

Ph: 0800 333 306;

Email: support@fertilitynz.org.nz

FertilityNZ is New Zealand's national network for those seeking support, information and news on fertility problems. It provides various services including:

- regional support and contact groups
- general advice and contact service
- comprehensive information brochures
- a forum for confidential feedback on any issues or concerns
- a chatroom where you can seek on-line support from people in similar situations.

Endometriosis New Zealand

www.nzendo.org.nz

Ph: +64 3 379 7959 (phone support line);

Email: info@nzendo.org.nz

Endometriosis New Zealand promotes awareness of endometriosis, provides information, education and raises funds to support endometriosis related initiatives. It includes disease information specifically designed for teenagers, a support group network, regular seminars and workshops and a free phone support line.

SANDS New Zealand

www.sands.org.nz

Ph: 027 44 91 019;

Email: info@sands.org.nz

SANDS is a self-help support group comprised of parents who have experienced the death of a baby through miscarriage, stillbirth, or shortly after birth. It provides 24-hour telephone support, information resources, monthly support meetings, name-giving certificates and other support.

REFERENCES

- 1 Assisted Conception Taskforce (ACT). Trying to have a baby. Your step-by-step guide to assisted conception. Available at http://www.assistedconception.net/resources/act_ratgeber.pdf. Accessed: 05 March 2020.
- 2 Allahbadia GN. *J Obstet Gynaecol India* 2017;67(6):385–92.
- 3 Speroff L *et al*. Clinical Gynecologic Endocrinology and Infertility. 6th ed. 1999.
- 4 Gardner DK *et al* (Eds.). Textbook of Assisted Reproductive Techniques. 2nd ed. 2004.
- 5 The ESHRE Capri Workshop Group. *Hum Reprod Update* 2009;15(3):265–77.
- 6 Yang S *et al*. *Chin Med J (Engl)*. 2016; 129(23):2873–75.

Looking for more information?

Other booklets in the *Pathways to Parenthood* series are available at www.fertilityportal.com.au/merck:

- Your step by step guide to treating infertility
- Overcoming male infertility
- Female infertility & assisted reproductive technology (ART)
- Endometriosis
- Ovulation induction (OI)
- Intrauterine insemination (IUI)
- *In vitro* fertilisation (IVF) & intra-cytoplasmic sperm injection (ICSI)
- Managing the stress of infertility
- Creating families for same-sex couples